


Microbes of Potential Concern in Distribution System Biofilms

Kellogg J. Schwab Ph.D.

Johns Hopkins Bloomberg School of Public Health
Department of Environmental Health Sciences



Microorganisms of public health concern in Biofilms

- It is not just bacteria
 - Viruses, protozoa and fungi can contribute to morbidity and mortality
- Immune status is important
 - Immunocompromised individuals have increased susceptibility to infection
 - An opportunistic pathogen is one that usually causes disease only in those whose immune system is compromised

Immune status


- A weakened immune system may be due to:
 - Very young or old age
 - By 2030, nearly 20% of the total U.S. population will be over 65 years of age
 - Cancer therapy
 - the number of persons living with cancer has tripled
 - 3.0 million in 1971
 - 9.8 million in 2001
 - Many are on immunosuppressive medications
 - Pregnancy – 6 million
 - Chronic illness
 - Diabetes – 16 million (~6 % of US population)
 - Cardiovascular disease – 60 million (~22% of US population)
 - Human immunodeficiency virus
 - Over 400,000 individuals in the US are living with AIDS



Bacteria of Health Concern

- Opportunistic bacterial pathogens
 - *Legionella spp.*
 - *Mycobacteria spp.*
 - *Pseudomonas spp.*
 - *Aeromonas spp.*
- Primary bacterial pathogens
 - *Campylobacter spp.*
 - Enterohemorrhagic *E. coli*
 - *Salmonella typhimurium*
 - *Shigella*
 - *Yersinia*
 - *Helicobacter pylori*

Bacteria in Biofilms



<http://www3.niaid.nih.gov/labs/aboutlabs/lhbp/pathogenMolecularGeneticsSection/>

- Biofilms are remarkably difficult to treat with antimicrobials.
 - Antimicrobials may be readily inactivated or fail to penetrate into the biofilm
 - bacteria within biofilms have increased (up to 1000-fold higher) resistance to antimicrobial compounds
 - even though these same bacteria are sensitive to these agents if grown under planktonic conditions
- Biofilms increase the opportunity for gene transfer between/among bacteria
- Certain species of bacteria communicate with each other within the biofilm.
 - As their density increases, the organisms secrete low molecular weight molecules that signal when the population has reached a critical threshold. This process, called quorum sensing, is responsible for the expression of virulence factors
- Bacteria express new, and sometimes more virulent phenotypes when growing within a biofilm.
- Bacteria embedded within biofilms are resistant to both immunological and non-specific defense mechanisms of the body

Legionella – An important biofilm bacteria

- At least 46 species and 70 serogroups have been identified.
 - *Legionella pneumophila*, an ubiquitous aquatic Gram-negative bacteria that thrives in warm environments (32°- 45°C)
 - causes over 90% of legionnaires' disease (LD) in the United States.
- LD is the more severe form of legionellosis
 - characterized by pneumonia commencing 2-10 days after exposure.
- Pontiac fever is an acute-onset, flu-like, non-pneumonic illness, occurring within a few hours to two days of exposure.



http://www.nrc-cnrc.gc.ca/education/bio/gallery/legionella_e.html



Legionella Transmission

- Person-to-person transmission has not been reported to occur
- Inhalation of contaminated aerosols
 - cooling towers,
 - showers and faucets
 - aspiration of contaminated water
- Legionnaire's disease is an acute respiratory infection which can cause a broad spectrum of symptoms from mild cough and fever to a serious pneumonia

Legionella Incidence

- An estimated 8,000-18,000 cases of *Legionella* occur each year in the United States, but because of under-diagnosis and underreporting, only 2 to 10% of estimated cases are identified.
 - Most legionnaires' disease cases are sporadic;
 - 20-25% are nosocomial
 - 10%-20% can be linked to outbreaks.
 - Pontiac fever has been recognized only during outbreaks.
- Death occurs in 10%-15% of legionnaires' disease cases:
 - a substantially higher proportion of fatal cases occur during nosocomial outbreaks.
 - Pontiac fever is a self-limited disease that requires no treatment.

Free-living Protozoa

- Amoebae and Acanthamoebae
 - Intracellular replication of *L. pneumophila* within protozoa play a major role in the transmission of Legionnaires' disease.
 - Protozoa provide habitats for *Legionella* environmental survival and replication
 - Other opportunistic pathogens can also be associated with free-living protozoa
 - *Mycobacterium* spp., *Pseudomonas* spp., *Burkholderia* spp., *Aeromonas* spp., *H. pylori* etc

Mycobacterium avium Complex (MAC)

- MAC consists of 28 serovars of two distinct species:
 - *Mycobacterium avium*
 - *Mycobacterium intracellulare*
- Exposure to MAC organisms may occur through
 - contact with or ingestion, aspiration, or aerosolization of potable water containing the organisms.
 - inhalation of air with contaminated soil particles
 - consumption of contaminated food




Mycobacterium avium Complex

- The clinical presentation of MAC infections can include a productive cough, fatigue, fever, weight loss, and night sweats
- The symptoms encountered with MAC infections result from colonization of either the respiratory or the gastrointestinal tract, with possible dissemination to other locations in the body.

Mycobacterium avium Complex

- Disseminated infections are usually associated with HIV infection.
 - Incidence is decreasing among HIV- infected patients as a result of new treatment modalities
 - combination therapy with nucleoside reverse transcriptase inhibitors and protease inhibitors
 - antimycobacterial prophylaxis



<http://www.md.huji.ac.il/mirror/webpath/AIDS039.jpg>

MAC involving spleen



Mycobacterium avium Complex

- *M. avium* can infect individuals with chronic lung conditions
 - bronchiectasis, emphysema and cystic fibrosis
- Lung disease secondary to aerosolized MAC from hot tubs has recently been described in young, immunocompetent individuals.



Mycobacterium avium Complex

- Unlike gastrointestinal pathogens, where fecal indicators such as *E. coli* can be used to indicate their potential presence, no suitable indicators have been identified to signal increasing concentrations of MAC organisms in water systems.

Pseudomonas spp



© 2004 Dennis Kunkel Microscopy, Inc.

<http://www.ehagroup.com/epidemiology/illnesses/images/pseudomonas-aeruginosa.jpg>

- *Pseudomonas aeruginosa* is common in soil, water, and vegetation.
- It is found on the skin of healthy individuals and has been isolated from the throat (5%) and stool (3%) of nonhospitalized patients.
- The gastrointestinal carriage rates increase in hospitalized patients to 20% within 72 hours of admission.

Pseudomonas spp.

- *Pseudomonas aeruginosa* is an opportunistic pathogen
 - respiratory system infections
 - urinary tract infections,
 - Dermatitis & soft tissue infections
 - bacteremia
 - bone and joint infections
 - gastrointestinal infections
- *Pseudomonas aeruginosa* infection is a serious problem in patients hospitalized with cancer, cystic fibrosis, and burns. The case fatality rate in these patients is 50%
- Many *Pseudomona spp.* are resistant to multiple antibiotics




<http://www.microbes-edu.org/etudiant/imgbgn/pyomain.jpg>

Aeromonas spp.

- The species principally associated with gastroenteritis are
 - *A. caviae*, *A. hydrophila*, and *A. veronii*
- Aeromonads have been commonly isolated from patients with gastroenteritis although their role in disease causation remains unclear.
 - They are also associated with
 - sepsis and wounds
 - eye, respiratory tract, and other systemic infections
- The pathogenesis of *Aeromonas* infections remains poorly understood
 - They express a range of virulence factors, including attachment mechanisms and produce a number of toxins.

Primary Bacterial Pathogens

- *Many enteric bacteria have the potential to attach to biofilms where they can aggregate and be protected from disinfection.*
 - *Campylobacter spp.*
 - Enterohemorrhagic *E. coli*
 - *Salmonella typhimurium*
 - *Shigella*
 - *Yersinia*
 - *Helicobacter pylori*





Enteric Viruses



- Transmitted via fecal-oral route – predominantly from human waste
- Viruses cannot replicate outside of specific hosts thus they can accumulate but not grow in a biofilm
 - Protection from disinfectants
 - Aggregation of virus
- Enteric viruses are non-enveloped and thus resistant to environmental degradation and chemical inactivation
 - Persist for weeks to months in cool aqueous environments
- The infectious dose for most human enteric viruses is very low (10-100 particles) and thus even very low levels are of public health concern
- There is minimal to no correlation between bacterial indicators and viruses in potable water



Enteric Viruses of Public Health Concern

- Calicivirus
 - Norovirus and Sapovirus
- Hepatitis A virus
- Adenovirus
- Rotavirus
- Coxsackievirus
- Echovirus
- Astrovirus


Norovirus (NoV)




- A genus in *Caliciviridae* family
 - Genetically diverse >150 strains
- The single most important cause of nonbacterial gastroenteritis in US and throughout the world
 - 23 million cases per year in US
- Small round structure, 27-30 *nm* dia.
- Non-enveloped protein capsid, positive sense single strand RNA genome
- Causes self-limiting intestinal illness
 - Major concern in nosocomial settings

Mimivirus

- **Mimivirus**
(*Acanthamoeba polyphaga*)
 - Mimivirus may be a causative agent of some forms of pneumonia



<http://en.wikipedia.org/wiki/Image:Mimivirus.jpg>



<http://www.stanford.edu/group/virus/mimi/2005/mimipic1.jpg>



Primary Protozoa

- Like viruses most primary protozoal pathogens cannot replicate outside of specific hosts thus they can accumulate but not grow in a biofilm
 - Present in water in a non-reproductive protective stage (e.g. cyst, oocyst)
- Human pathogenic protozoa are resistant to environmental degradation and chemical inactivation
 - Persist for months in cool aqueous environments
- The infectious dose for most human enteric protozoa is very low (10s-100s of (oo)cysts) and thus even very low levels are of public health concern
- There is minimal to no correlation between bacterial indicators and protozoa in potable water




Protozoa of Public Health Concern

- *Cryptosporidium*
- *Giardia*
- *Microsporidia*
- *Toxoplasma gondii*
- *Cyclospora*
- *Entamoeba*
- *Acanthamoeba*
- *Naegleria*

Cryptosporidium

- Oocysts are very infectious
 - Infectious dose 10s to 100s
- 3-11 day incubation (average is 7 days)
- May result in asymptomatic carriage
- Self-limiting in healthy individuals
 - Watery diarrhea without blood
 - Illness lasts 10-14 days
- Immunocompromised patients
 - >50 stools/day with tremendous fluid loss
 - Can be severe and last for months
 - 10-15% of AIDS patients die of complications related to cryptosporidiosis






Fungi

- Fungi are classified as
 - Molds – branching, threadlike filaments
 - Yeasts – single-celled organisms that reproduce by budding
- Fungi are ubiquitous in the environment and have been found in water distribution systems
 - Opportunistic and infrequently cause illness
 - Several spp. can produce toxins

Fungi

- *Aspergillus* spp.
 - *A. fumigatus*; *A. flavus*; *A. niger*
 - Pulmonary disease
- *Cryptococcus neoformans*
 - Lung infections, meningitis
- *Candida albicans*
 - Vaginal, urinary and esophageal infections
- *Stachybotrys chartarum*
 - Pulmonary disease



<http://www.pall.com/images/flavcf.gif>



Detection Strategies

- Many bacteria can enter a viable but not cultivable (VBNC) state making culture analysis challenging
- Culture analysis reflective of infectivity
 - Risk to public health
- Molecular techniques are sensitive and specific but most do not provide information on the infectious nature of the detected microorganism

Control of Microorganisms of Public Health Concern in Biofilms

- The number of immunosuppressed individuals is increasing
 - Increased health risks due to opportunistic pathogens
- Microorganisms other than bacteria are important public health risks
 - Viruses, protozoa, fungi
- Control of microbial contamination requires maintenance of a constant disinfectant residual throughout the water system.
- Treatment strategies targeting free-living amoebae should lead to improved control of *L. pneumophila* and other pathogens.
- Biofilms are unique environments that will continue to be a challenge for the water industry